Jap. 8814/64, clg. (Non-Con.) 18.5.60 (Jap.) as 35704/60.  Pub. 28.5.64. Yamanouchi Pharm. Co., Ltd.  NEW  The following process and cmpds. of the general formula (1)  CH <sub>2</sub> =CH-CH <sub>2</sub> (CH <sub>2</sub> ) <sub>4</sub> COOR  CH <sub>3</sub> -CH <sub>3</sub> -CH-(CH <sub>2</sub> ) <sub>4</sub> COOR  HI  CH <sub>4</sub> -CH <sub>3</sub> -CH-(CH <sub>2</sub> ) <sub>4</sub> COOR  NH <sub>2</sub> CH <sub>2</sub> -CH-(CH <sub>2</sub> ) <sub>4</sub> COOR  CC  CC  CC  CC  CC  CC  CC  CC  CC	12465
(II) is intermediate for prepa-blook (II) is intermediate for prepa-blook (II) is intermediate for prepa-blook (II) is intermediate for preparation of blook (II) was reflex (II) ester (II) was reflex soln. made alkaline with 25% N40 mins. The mixture was manydrochloric acid, extracted wiether evap. and product distillicapto-octanoic acid (9.5 g.) b. 150°/0.05 mm.	
of medicines (e.g.  - \( \times \) -	8.5.60 YAMA